



NORTH-HOLLAND

Operator Versions of Some Classical Inequalities

B. Mond

School of Mathematics

La Trobe University

Bundoora 3083, Victoria, Australia

J. Pečarić

Faculty of Textile Technology

University of Zagreb

Zagreb, Croatia

J. Šunde

Applied Mathematics Department

The University of Adelaide

Adelaide, South Australia 5005, Australia

and

S. Varošanec

Department of Mathematics

University of Zagreb

Zagreb 10,000 Croatia

Submitted by Simo Puntanen

ABSTRACT

Using the theory of connections developed in a paper by Kubo and Ando, operator versions of many classical inequalities are obtained. Finally, it is shown how abstract solidarities, an extension of the Kubo-Ando theory, can be used to obtain operator inequalities. © 1997 Elsevier Science Inc.

LINEAR ALGEBRA AND ITS APPLICATIONS 264:117–126 (1997)

© 1997 Elsevier Science Inc. All rights reserved.
655 Avenue of the Americas, New York, NY 10010

0024-3795/97/\$17.00
PII S0024-3795(97)00234-6

1. INTRODUCTION

Let us consider bounded, linear, and positive (i.e. positive semidefinite) operators on an infinite dimensional Hilbert space. A scalar multiple of the identity operator is denoted by the scalar itself; in particular, 1 is the identity operator. The order relation $A \leq B$ means that $B - A$ is positive. That $A_1 \geq A_2 \geq \dots$ and A_n converges strongly to A is denoted by $A_n \downarrow A$.

A binary operator σ on the class of positive operators, $(A, B) \rightarrow A \sigma B$, is called a *connection* if the following requirements are fulfilled [1]:

- (I) $A \leq C$ and $B \leq D \Rightarrow A \sigma B \leq C \sigma D$,
- (II) $C(A \sigma B)C \leq (CAC) \sigma (CBC)$,
- (III) $A_n \downarrow A$ and $B_n \downarrow B \Rightarrow (A_n \sigma B_n) \downarrow A \sigma B$.

A *mean* is a connection with normalization condition

- (IV) $1 \sigma 1 = 1$.

The following results are also valid [1]: Every mean σ possesses the property

- (IV') $A \sigma A = A$ for every A .

Every connection σ possesses the property

- (I') $(A \sigma B) + (C \sigma D) \leq (A + C) \sigma (B + D)$

The simplest examples of means are

Arithmetic mean: $A \nabla B = \frac{1}{2}(A + B)$.

Harmonic mean: $A ! B = 2(A^{-1} + B^{-1})^{-1}$.

Geometric mean: $A \# B = A^{1/2}(A^{-1/2}BA^{-1/2})^{1/2}A^{1/2}$ for invertible A and B .

Moreover, weighted versions of these means can also be defined.

Let A, B be invertible and $\lambda \in (0, 1)$ be a real number. Then the arithmetic, geometric, and harmonic means are defined, respectively, by

$$\begin{aligned} A \nabla_{\lambda} B &= \lambda A + (1 - \lambda) B, \\ A \#_{\lambda} B &= A^{1/2} (A^{-1/2} B A^{-1/2})^{1-\lambda} A^{1/2}, \\ A !_{\lambda} B &= [\lambda A^{-1} + (1 - \lambda) B^{-1}]^{-1}. \end{aligned}$$

It is well known that the following inequalities are valid:

$$A !_{\lambda} B \leq A \#_{\lambda} B \leq A \nabla_{\lambda} B. \quad (1.1)$$

Of course a connection (or a mean) σ is symmetric if

$$A \sigma B = B \sigma A.$$

The following result is valid [2]: Let $A_1 \leq \dots \leq A_n$ (not all equal) and $B_1 \leq \dots \leq B_n$ (not all equal) be bounded linear, positive, and invertible operators such that $A_1 = B_1$ and $A_n = B_n$, and let $a_1 \leq \dots \leq a_n$ be positive numbers. If σ is a symmetric mean, then

$$\left(\sum_{i=1}^{n-1} a_i \Delta A_i \right) \sigma \left(\sum_{i=1}^{n-1} a_i \Delta B_i \right) \leq \sum_{i=1}^{n-1} a_i \Delta (A_i \sigma B_i), \quad (1.2)$$

where Δ is a finite difference, i.e., $\Delta a_i = a_{i+1} - a_i$, $i = 1, \dots, n-1$.

In fact, in [2], we have proved a more general result for two symmetric means.

In this paper, we shall show that (1.2) (indeed, a more general result) and operator versions of some general inequalities are consequences of (I'). Finally, it is shown how abstract solidarities, an extension of the Kubo-Ando theory, can be used to obtain operator inequalities.

2. OPERATOR VERSIONS OF CAUCHY, HÖLDER, AND OTHER CLASSICAL INEQUALITIES

Mathematical induction gives, from (I'):

THEOREM 2.1. *Let A_i, B_i , $i = 1, \dots, n$, be bounded linear, and positive operators, and let σ be a connection. Then*

$$\sum_{i=1}^n (A_i \sigma B_i) \leq \left(\sum_{i=1}^n A_i \right) \sigma \left(\sum_{i=1}^n B_i \right). \quad (2.1)$$

In the next examples, we assume that A_i and B_i are invertible.

EXAMPLES.

(1) Cauchy's inequality:

$$\sum_{i=1}^n A_i^2 \# B_i^2 \leq \left(\sum_{i=1}^n A_i^2 \right) \# \left(\sum_{i=1}^n B_i^2 \right). \quad (2.2)$$

(2) Hölder's inequality: Let $p, q > 0$ with $p^{-1} + q^{-1} = 1$. Then

$$\sum_{i=1}^n A_i^p \#_{1/p} B_i^q \leq \left(\sum_{i=1}^n A_i^p \right) \#_{1/p} \left(\sum_{i=1}^n B_i^q \right). \quad (2.3)$$

(3) Minkowski's inequality:

$$\sum_{i=1}^n (A_i + B_i)^{-1} \leq \left[\left(\sum_{i=1}^n A_i^{-1} \right)^{-1} + \left(\sum_{i=1}^n B_i^{-1} \right)^{-1} \right]^{-1} \quad (2.4)$$

Indeed, this last inequality follows by letting σ be the parallel sum, i.e.,

$$A \sigma B = (A^{-1} + B^{-1})^{-1} \quad \text{and} \quad A_i = A_i^{-1}, \quad B_i = B_i^{-1}.$$

THEOREM 2.2. *Let $A_i, B_i, i = 1, \dots, n$, be bounded, linear, and positive operators such that*

$$A_1 - A_2 - \dots - A_n \geq 0 \quad \text{and} \quad B_1 - B_2 - \dots - B_n \geq 0. \quad (2.5)$$

Then

$$A_1 \sigma B_1 - \sum_{i=2}^n A_i \sigma B_i \geq \left(A_1 - \sum_{i=2}^n A_i \right) \sigma \left(B_1 - \sum_{i=2}^n B_i \right). \quad (2.6)$$

Proof. By the substitutions

$$A_1 \rightarrow A_1 - A_2 - \dots - A_n, \quad B_1 \rightarrow B_1 - B_2 - \dots - B_n$$

(2.1) becomes

$$(A_1 - A_2 - \dots - A_n) \sigma (B_1 - B_2 - \dots - B_n) + \sum_{i=2}^n A_i \sigma B_i \leq A_1 \sigma B_1,$$

i.e. (2.6). ■

In the following examples A_i and B_i are again invertible.

EXAMPLES.

(4) Aczél's inequality: If

$$A_1^2 - A_2^2 - \dots - A_n^2 > 0 \quad \text{and} \quad B_1^2 - B_2^2 - \dots - B_n^2 > 0,$$

then

$$A_1^2 \# B_1^2 - \sum_{i=2}^n A_i^2 \# B_i^2 \geq \left(A_1^2 - \sum_{i=2}^n A_i^2 \right) \# \left(B_1^2 - \sum_{i=2}^n B_i^2 \right). \quad (2.7)$$

(5) Popoviciu's inequality: If $p, q > 0$, $p^{-1} + q^{-1} = 1$, and

$$A_1^p - A_2^p - \cdots - A_n^p > 0, \quad B_1^q - B_2^q - \cdots - B_n^q > 0,$$

then

$$A_1^p \#_{1/p} B_1^q - \sum_{i=2}^n A_i^p \#_{1/p} B_i^q \geq \left(A_1^p - \sum_{i=2}^n A_i^p \right) \#_{1/p} \left(B_1^q - \sum_{i=2}^n B_i^q \right). \quad (2.8)$$

(6) Bellman's inequality: If

$$A_1^{-1} - A_2^{-1} - \cdots - A_n^{-1} > 0 \quad \text{and} \quad B_1^{-1} - B_2^{-1} - \cdots - B_n^{-1} > 0,$$

then

$$\begin{aligned} & (A_1 + B_1)^{-1} - \sum_{i=2}^n (A_i + B_i)^{-1} \\ & \geq \left[\left(A_1^{-1} - \sum_{i=2}^n A_i^{-1} \right)^{-1} + \left(B_1^{-1} - \sum_{i=2}^n B_i^{-1} \right)^{-1} \right]^{-1} \end{aligned} \quad (2.9)$$

REMARKS.

(1) Note that (2.4) and (2.9) can be given in the following forms:

$$\left(\sum_{i=1}^n (A_i + B_i)^{-1} \right)^{-1} \geq \left(\sum_{i=1}^n A_i^{-1} \right)^{-1} + \left(\sum_{i=1}^n B_i^{-1} \right)^{-1} \quad (2.4')$$

and

$$\begin{aligned} & \left((A_1 + B_1)^{-1} - \sum_{i=2}^n (A_i + B_i)^{-1} \right)^{-1} \\ & \leq \left(A_1^{-1} - \sum_{i=2}^n A_i^{-1} \right)^{-1} + \left(B_1^{-1} - \sum_{i=2}^n B_i^{-1} \right)^{-1} \end{aligned} \quad (2.9')$$

Note that the following generalization of (2.4') is obtained in [3] for positive invertible operators A_{ij} ($i = 1, \dots, n$, $j = 1, \dots, m$):

$$\sum_{j=1}^m \left(\sum_{i=1}^n A_{ij}^{-1} \right)^{-1} \leq \left[\sum_{i=1}^n \left(\sum_{j=1}^m A_{ij} \right)^{-1} \right]^{-1}. \quad (2.10)$$

We can use (2.10) in the proof of the following extension of (2.9'): If positive invertible operators A_{ij} ($i = 1, \dots, m$, $j = 1, \dots, n$) satisfy the conditions

$$A_{1j}^{-1} - A_{2j}^{-1} - \dots - A_{mj}^{-1} > 0, \quad j = 1, \dots, n,$$

then

$$\left(\sum_{j=1}^m A_{1j} \right)^{-1} - \sum_{i=2}^m \left(\sum_{j=1}^n A_{ij} \right)^{-1} \geq \left[\sum_{j=1}^n \left(A_{1j}^{-1} - \sum_{i=2}^m A_{ij}^{-1} \right)^{-1} \right]^{-1} \quad (2.11)$$

(2) A simpler form of Hölder's inequality is

$$\sum_{i=1}^n A_i \#_{\alpha} B_i \leq \left(\sum_{i=1}^n A_i \right) \#_{\alpha} \left(\sum_{i=1}^n B_i \right), \quad (2.3')$$

where $0 \leq \alpha \leq 1$. Setting $A_i \rightarrow A_i^s$, $B_i \rightarrow A_i^r$ in (2.3'), we get

$$\begin{aligned} \sum_{i=1}^n A_i^{\alpha s + (1-\alpha)r} & \leq \left(\sum_{i=1}^n A_i^s \right) \#_{\alpha} \left(\sum_{i=1}^n A_i^r \right) \\ & \leq \alpha \left(\sum_{i=1}^n A_i^s \right) + (1-\alpha) \left(\sum_{i=1}^n A_i^r \right), \end{aligned}$$

where we have used (1.1). In fact, we have proved that the function

$$x \mapsto \sum_{i=1}^n A_i^x$$

is convex.

3. AN OPERATOR VERSION OF PÓLYA'S INEQUALITY

THEOREM 3.1. *Let $A_1 \leq \dots \leq A_n$ (not all equal) and $B_1 \leq \dots \leq B_n$ (not all equal) be bounded, linear, and positive operators, let $a = \{a_1, \dots, a_n\}$ be a nondecreasing positive n -tuple of real numbers, and let σ be a connection. Then*

$$\left(\sum_{i=1}^{n-1} a_i \Delta A_i \right) \sigma \left(\sum_{i=1}^{n-1} a_i \Delta B_i \right) \leq \sum_{i=1}^n a_i \Delta (A_i \sigma B_i). \quad (3.1)$$

If a is a nonincreasing positive n -tuple of real numbers and $A_1 = B_1 = 0$, then a reverse inequality in (3.1) holds.

Proof.

(1) We have

$$\begin{aligned} & \sum_{i=1}^{n-1} a_i \Delta (A_i \sigma B_i) \\ &= a_n (A_n \sigma B_n) - a_1 (A_1 \sigma B_1) - \sum_{i=2}^n (A_i \sigma B_i) \Delta a_{i-1} \\ &= a_n (A_n \sigma B_n) - a_1 (A_1 \sigma B_1) - \sum_{i=2}^n [(A_i \Delta a_{i-1}) \sigma (B_i \Delta a_{i-1})] \\ &\geq (a_n A_n) \sigma (a_n B_n) - (a_1 A_1) \sigma (a_1 B_1) \\ &\quad - \left(\sum_{i=2}^n A_i \Delta a_{i-1} \right) \sigma \left(\sum_{i=2}^n B_i \Delta a_{i-1} \right) \quad [\text{by (2.1)}] \end{aligned}$$

$$\begin{aligned}
&\geq \left(a_n A_n - a_1 A_1 - \sum_{i=2}^n A_i \Delta a_{i-1} \right) \sigma \left(a_n B_n - a_1 B_1 - \sum_{i=2}^n B_i \Delta a_{i-1} \right) \\
&\quad \text{[by (2.6) for three terms]} \\
&= \left(\sum_{i=1}^{n-1} a_i \Delta A_i \right) \sigma \left(\sum_{i=1}^{n-1} a_i \Delta B_i \right).
\end{aligned}$$

(2) Moreover if a is nonincreasing and $A_1 = B_1 = 0$, we have, since $-a$ is nondecreasing, as a consequence of (2.1),

$$\begin{aligned}
&\sum_{i=1}^{n-1} a_i \Delta (A_i \sigma B_i) \\
&= a_n (A_n \sigma B_n) + \sum_{i=2}^n (A_i \sigma B_i) \Delta (-a_{i-1}) \\
&= (a_n A_n) \sigma (a_n B_n) + \sum_{i=2}^n [A_i \Delta (-a_{i-1})] \sigma [B_i \Delta (-a_{i-1})] \\
&\leq (a_n A_n) \sigma (a_n B_n) + \left(\sum_{i=2}^n A_i \Delta (-a_{i-1}) \right) \sigma \left(\sum_{i=2}^n B_i \Delta (-a_{i-1}) \right) \\
&\leq \left(a_n A_n + \sum_{i=2}^n A_i \Delta (-a_{i-1}) \right) \sigma \left(a_n B_n + \sum_{i=2}^n B_i \Delta (-a_{i-1}) \right) \\
&= \left(\sum_{i=1}^{n-1} a_i \Delta A_i \right) \sigma \left(\sum_{i=1}^{n-1} a_i \Delta B_i \right). \quad \blacksquare
\end{aligned}$$

REMARK. Theorem 3.1 gives the inequality (1.2) for arbitrary connections (not only symmetric means) and also without the conditions $A_1 = B_1$ and $A_n = B_n$. Also, we have a converse result in our theorem.

COROLLARY 3.1. Let $A_1 \leq \dots \leq A_n$ (not all equal) and $B_1 \leq \dots \leq B_n$ (not all equal) be bounded, linear, positive, and invertible operators, and let $\lambda \in [0, 1]$. If a is a nondecreasing n -tuple of positive numbers, then

$$\left(\sum_{i=1}^{n-1} a_i \Delta A_i \right) \#_{\lambda} \left(\sum_{i=1}^{n-1} a_i \Delta B_i \right) \leq \sum_{i=1}^{n-1} a_i (A_i \#_{\lambda} B_i) \quad (3.2)$$

and

$$\left(\sum_{i=1}^{n-1} a_i \Delta A_i \right) !_{\lambda} \left(\sum_{i=1}^{n-1} a_i \Delta B_i \right) \leq \sum_{i=1}^{n-1} a_i (A_i !_{\lambda} B_i) \quad (3.3)$$

If a is a nonincreasing n -tuple of positive numbers and $A_1 = B_1 = 0$, then reverse inequalities in (3.2) and (3.3) are valid.

REMARKS. Inequalities (3.2) and (3.3) were proved in [2], but with the additional conditions $A_1 = B_1$ and $A_n = B_n$.

4. INEQUALITIES FOR SOLIDARITIES

An extension of the Kubo-Ando theory was given by J. I. Fujii, M. Fujii, and Y. Seo [4].

A binary operation s on positive operators is an *abstract solidarity* if it satisfies, assuming the existence of $A s B$ as a bounded operator, that

- (S1) $B \leq C$ implies $A s B \leq A s C$,
- (S2r) $B_n \downarrow B$ implies $A s B_n \downarrow A s B$,
- (S2ℓ) $A_n \rightarrow A$ strongly implies $A_n s 1 \rightarrow A s 1$ strongly, and
- (S3) $T^*(A s B)T \leq T^*AT s T^*BT$.

The solidarity s is superadditive in that

$$(A + B) s (C + D) \geq A s C + B s D. \quad (4.1)$$

Of special interest is the relative operator entropy $S(A|B)$ for invertible A, B defined by

$$S(A|B) = A^{1/2}(\log A^{-1/2}BA^{-1/2})A^{1/2}$$

Of course, using (4.1), we can prove the following by mathematical induction: Let $A_i, B_i, i = 1, \dots, n$, be positive operators; then

$$\sum_{i=1}^n (A_i s B_i) \leq \left(\sum_{i=1}^n A_i \right) s \left(\sum_{i=1}^n B_i \right). \quad (4.2)$$

Also, if (2.5) holds, then

$$A_1 s B_1 - \sum_{i=2}^n A_i s B_i \geq \left(A_1 - \sum_{i=2}^n A_i \right) s \left(B_1 - \sum_{i=2}^n B_i \right). \quad (4.3)$$

For the operator entropy, we have

$$\sum_{i=1}^n S(A_i|B_i) \leq S\left(\sum_{i=1}^n A_i \middle| \sum_{i=1}^n B_i\right) \quad (4.4)$$

and, if (2.5) holds,

$$S(A_1|B_1) - \sum_{i=2}^n S(A_i|B_i) \geq S\left(A_1 - \sum_{i=2}^n A_i \middle| B_1 - \sum_{i=2}^n B_i\right). \quad (4.5)$$

As in Theorem 3.1, we can also prove

THEOREM 4.1. *Let $A_1 \leq \dots \leq A_n$ (not all equal) and $B_1 \leq \dots \leq B_n$ (not all equal) be bounded, linear, and positive operators, let $a = \{a_1, \dots, a_n\}$ be a nondecreasing n -tuple of positive numbers, and let s be an abstract solidarity. Then*

$$\left(\sum_{i=1}^{n-1} a_i \Delta A_i \right) s \left(\sum_{i=1}^n a_i \Delta B_i \right) \leq \sum_{i=1}^{n-1} a_i \Delta (A_i s B_i). \quad (4.6)$$

If a is a nonincreasing n -tuple of positive numbers and $A_1 = B_1 = 0$, then a reverse inequality in (4.6) is valid.

In the case of operator entropy, (4.6) becomes

$$S\left(\sum_{i=1}^{n-1} a_i \Delta A_i \middle| \sum_{i=1}^{n-1} a_i \Delta B_i\right) \leq \sum_{i=1}^{n-1} a_i S(A_i|B_i). \quad (4.7)$$

REFERENCES

- 1 F. Kubo and T. Ando, Means of positive linear operators, *Math. Ann.* 246:205–224 (1980).
- 2 B. Mond, J. Pečarić, J. Šunde, and S. Varošanec, Pólya's inequality for positive linear operators, *Rad Hazu*, to appear.
- 3 W. N. Anderson, Jr., and R. J. Duffin, Series and parallel addition of matrices, *J. Math. Anal. Appl.* 26:576–594 (1969).
- 4 J. I. Fujii, M. Fujii, and Y. Seo, An extension of the Kubo-Ando theory: Solidarities, *Math. Japon.* 35:387–396 (1990).

Received 16 May 1996; final manuscript accepted 16 December 1996